


CATEGORICAL q -DEFORMED RATIONAL
NUMBERS VIA BRIDGELAND STABILITY
CONDITIONS


Asilata Bapat (ANU)

+ Louis Becker,
Anthony Licata

Outline



Outline



Fractional linear action of B_3

$$B_3 = \langle \sigma_1, \sigma_2 \mid \sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2 \rangle$$

There is a homomorphism

$$B_3 \rightarrow PSL_2(\mathbb{Z}) :$$

$$\sigma_1 \mapsto \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}, \quad \sigma_2 \mapsto \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$$

Fractional linear action of B_3

$PSL_2(\mathbb{Z})$ acts on $\mathbb{R} \cup \{\infty\}$ via fractional linear transformations:

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \cdot \left(\frac{r}{s} \right) := \frac{ar + bs}{cr + ds}$$

- * B_3 acts on $\mathbb{R} \cup \{\infty\}$.
- * The action preserves $\mathbb{Q} \cup \{\infty\}$.

Fractional linear action of B_3

Can be realised via continued fractions.

Let $\frac{\gamma}{s} = a_1 + \cfrac{1}{a_2 + \cfrac{1}{\dots + a_{2n}}}$

Then

$$\frac{\gamma}{s} = \sigma_1^{-a_1} \sigma_2^{a_2} \sigma_1^{-a_3} \sigma_2^{a_4} \dots \sigma_1^{-a_{2n-1}} \sigma_2^{a_{2n}}(\infty)$$

Classical (right) q -deformed rationals

Consider deformed matrices :

$$\sigma_{1,q} := \begin{bmatrix} q^1 & -q^{-1} \\ 0 & 1 \end{bmatrix}, \quad \sigma_{2,q} := \begin{bmatrix} 1 & 0 \\ 1 & q^{-1} \end{bmatrix}$$

These generate a copy of B_3 in

$$\mathrm{PSL}_2(\mathbb{Z}[q^\pm]).$$

Classical (right) q -deformed rationals

Let $\frac{r}{s} = a_1 + \frac{1}{a_2 + \frac{1}{\dots + a_{2n}}}$

The corresponding right q -deformation is:

$$\left[\frac{r}{s} \right]_q^{\#} = \sigma_{1,q}^{-a_1} \sigma_{2,q}^{a_2} \cdots \sigma_{1,q}^{-a_{2n-1}} \sigma_{2,q}^{a_{2n}} (\infty)$$

[Monier-Genoud - Ovsienko]


Left q -deformed rationals

Let $\frac{r}{s} = a_1 + \frac{1}{a_2 + \frac{1}{\dots + a_{2n}}}$

The corresponding left q -deformation is:

$$\left[\frac{r}{s} \right]_q^b = \sigma_{1,q}^{-a_1} \sigma_{2,q}^{a_2} \cdots \sigma_{1,q}^{-a_{2n-1}} \sigma_{2,q}^{a_{2n}} \left(\frac{1}{1-q} \right).$$

Outline



Categorical interlude

\mathcal{C} = 2-CY category of connected graph Γ

[categorifies Burau rep of B_r]

Main example for this talk:

$$\Gamma = \text{---} \bullet$$

(A_2 graph)


$$\& B_r = B_3$$

Categorical interlude

\mathcal{C} = 2-CY category of connected graph Γ

[categorifies Burau rep of B_r]


Important features:

- $\mathcal{C} = \langle P_i \mid i \text{ vertex} \rangle$ 
- Can be realised via dg modules over the zigzag algebra of Γ .

Categorical Br action


The objects P_1 & P_2 are
"spherical":

$$\text{End}^m(P_i) = \begin{cases} \mathbb{C} & \text{if } m = 0, 2 \\ 0 & \text{otherwise} \end{cases}$$



Categorical Br action

The objects P_1 & P_2 are "spherical":




$$\text{End}^m(P_i) = \begin{cases} \mathbb{C} & \text{if } m = 0, 2 \\ 0 & \text{otherwise} \end{cases}$$

Any spherical object x defines autoequivalence $\sigma_x : \mathcal{C} \xrightarrow{\sim} \mathcal{C}$.

In particular, $\sigma_{P_1}, \sigma_{P_2} : \mathcal{C} \xrightarrow{\sim} \mathcal{C}$.

Categorical Br action

The functors σ_{P_1} & σ_{P_2} braid:



$$\sigma_{P_1} \sigma_{P_2} \sigma_{P_1} \simeq \sigma_{P_2} \sigma_{P_1} \sigma_{P_2}$$

Therefore we have a (weak) action

of B_3 on \mathcal{C}_{A_2} with

$$\sigma_1 \mapsto \sigma_{P_1} \text{ & } \sigma_2 \mapsto \sigma_{P_2}.$$

Bridgeland stability conditions & Br-action

We will encounter q-rationals again by taking "limiting q-sizes" of objects in \mathcal{C} .

These are provided by Bridgeland stability conditions.

Bridgeland stability conditions & Br-action

A stability condition τ is data on \mathcal{C} that yields a family of metrics on \mathcal{C} : each arrow in \mathcal{C} has a length

The size of $X \in \text{ob } \mathcal{C}$ is measured by "pulling tight" to a geodesic path

$$0 \rightarrow X_1 \rightarrow X_2 \rightarrow X_3 \rightarrow X$$

with "semistable" segments.


Bridgeland stability conditions & Br-action

The size of $X \in \text{ob } \mathcal{C}$ is measured by "pulling tight" to a geodesic $0 \rightarrow X$.

This is called the "q-mass" of X wrt τ .

If

$$X =$$



, then

$$m_{q,\tau}(X) = \sum q^{\phi(A_i)} \cdot |A_i|$$

Bridgeland stability conditions & B_r -action

[Bridgeland] $\text{Stab } \mathcal{C}$ is a complex manifold.

Since $B_r \subset \mathcal{C}$, we also have

$$B_r \subset \text{Stab } \mathcal{C}.$$

Bridgeland stability conditions & B_Γ -action

[Bridgeland] $\text{Stab } \mathcal{C}$ is a complex manifold.

When $\Gamma = A_2$, $\text{Stab } \mathcal{C}$ (modulo \mathbb{C} -action) is homeomorphic to the upper half plane.


Under this correspondence, B_3 acts by fractional linear maps via $\text{PSL}_2(\mathbb{Z})$.

Limiting operations on Stab 4

Limiting operations on $\text{Stab } \mathcal{C}$

① Fix $\beta \in B_r$ and $\tau \in \text{Stab } \mathcal{C}$.


Consider $\lim_{n \rightarrow \infty} \beta^n \tau$.



Limiting operations on $\text{Stab } \mathcal{C}$

① Fix $\beta \in B_r$ and $\tau \in \text{Stab } \mathcal{C}$.

Consider $\lim_{n \rightarrow \infty} \beta^n \tau$.




[BDL] Taking $\beta = \delta_x$ for x spherical :

$$\lim_{n \rightarrow \infty} m_{\beta^n \tau, q} (Y) = q\text{-dim Hom}(X, Y)$$

up to simultaneous scalar

Limiting operations on Stab 4


②




Shrink all but
one of the simple
semistables to zero

Limiting operations on Stab 4

②



Shrink all but
one of the simple
semistables to zero



In the limit, the
q-mass counts the
“q-occurrences” of the
remaining semistable
in any given object.

Limiting operations on Stab \mathbb{C}

Moral : Limits may not make sense as stability conditions, but their q-masses make sense.

Limiting operations on $\text{Stab } \mathcal{C}$

Moral : Limits may not make sense as stability conditions, but their q-masses make sense.

Mass map

$$\text{Stab } \mathcal{C} \xrightarrow{\quad} \mathbb{P}\mathbb{R}^S$$

$$\tau \longmapsto [x \mapsto m_{q,\tau}(x)]/\sim$$

Mass map & compactification

- [BDL, BBL] The mass map is injective, and $\overline{\text{Stab}}^g_{\mathcal{C}}$ is compact.


Mass map & compactification

- [BDL, BBL] The mass map is injective, and $\overline{\text{Stab}}^q_{\mathcal{C}}$ is compact.
- In the boundary, we see :


$$\overline{\text{hom}} := \lim_{n \rightarrow \infty} M_{\beta^n, q} \text{ for } \beta = \text{spherical twist}$$

$\text{occ} := q\text{-occurrences of a fixed semistable}$

Outline



The story of the 3-strand braid group




The story of the 3-strand braid group

$$B_3 = \langle \sigma_1, \sigma_2 \mid \sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2 \rangle$$

$$B_3 \rightarrow PSL_2(\mathbb{Z})$$

$$\sigma_1 \mapsto \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}, \quad \sigma_2 \mapsto \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$$




The story of the 3-strand braid group

$$B_3 = \langle \sigma_1, \sigma_2 \mid \sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2 \rangle$$

$$B_3 \rightarrow PSL_2(\mathbb{Z})$$

$$\sigma_1 \mapsto \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}, \quad \sigma_2 \mapsto \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$$



- $PSL_2(\mathbb{Z})$, and hence B_3 , acts on $\mathbb{C} \cup \{\infty\}$ by fractional linear transformations
- Action preserves \mathbb{H} and $\mathbb{R} \cup \{\infty\}$

The story of the 3-strand braid group

For the remainder of the talk, take

$$\mathcal{C} = \mathcal{C}(\bullet - \bullet) = \langle P_1, P_2 \rangle \circlearrowleft B_3$$

Fact :

$$\begin{array}{ccc} \text{Stab } \mathcal{C} & \simeq & \mathbb{H} \\ \mathcal{C} & & \mathcal{O} \\ B_3 & & B_3 \text{ via } \mathrm{PSL}_2(\mathbb{Z}) \end{array}$$

The story of the 3-strand braid group

Take $\mathcal{C} = \mathcal{C}(\bullet - \bullet) = \langle P_1, P_2 \rangle \wr B_3$

Thm [BDL]: For $q=1$:

- ① $\overline{\text{hom}}$ and occ coincide.
- ② $\overleftarrow{\text{hom}}_X \mapsto \pm \frac{\overline{\text{hom}}(X, P_2)}{\overline{\text{hom}}(X, P_1)}$ is a

B_3 -equivariant bijection from the spherical objects of \mathcal{C} to $\mathbb{Q} \cup \{\infty\}$


The $\overline{\text{hom}}$ functionals as rationals

At $q=1$: The rationals can be recovered as the quotients

$\pm \frac{\overline{\text{hom}}(X, P_2)}{\overline{\text{hom}}(X, P_1)}$ as X ranges over the spherical objects of \mathcal{C} .

The $\overline{\text{hom}}$ functionals as rationals

Pictorially, at $g=1$:



The q -deformed story for B_3

Question : Can we recover the q -rationals via some deformation of the quotients $\pm \frac{\hom(x, P_2)}{\hom(x, P_1)}$?

Answer : Yes, and more!

The q -deformed story for B_3


Thm [BBL]

① $\pm q^{(1)} \frac{\text{occ}(P_2, x)}{\text{occ}(P_1, x)}$ are exactly the classical
(right) q -deformed rationals of [M-G-O]

② $\pm q^{(1)} \frac{\overline{\text{hom}}(x, P_2)}{\overline{\text{hom}}(x, P_1)}$ is a new q -deformation
of \mathbb{Q} . These are
exactly the left q -rationals.

The q -deformed story for B_3

The right q -rationals at $q=1$:



The q -deformed story for B_3

Thm [cont'd]

③ $\overline{\hom}_X \mapsto \pm q^{\epsilon} \frac{\overline{\hom}_q(X, P_2)}{\overline{\hom}_q(X, P_1)}$ and

$$\text{occ}_X \mapsto \pm q^{\epsilon} \frac{\text{occ}(P_2, X)}{\text{occ}(P_1, X)}$$
 are B_3 -equivariant.

The q -deformed story for B_3

Thm [cont'd]

(3) $\overline{\text{hom}}_X \mapsto \pm q^{\epsilon} \frac{\overline{\text{hom}}_q(X, P_2)}{\overline{\text{hom}}_q(X, P_1)}$ and

$$\text{occ}_X \mapsto \pm q^{\epsilon} \frac{\text{occ}(P_2, X)}{\text{occ}(P_1, X)}$$
 are B_3 -equivariant.

The B_3 -action on the right is by fractional linear transformations via deformed B_3 matrices.

The q -deformed story for B_3

Upshot: for $\frac{r}{s} \in \mathbb{Q} \cup \{\infty\}$, we have

$$\textcircled{1} \quad \left[\frac{r}{s} \right]_q^{\#} = \pm q^{l'} \frac{\text{occ}(P_2, X)}{\text{occ}(P_1, X)} \quad \begin{matrix} \text{right } q\text{-deformed} \\ \text{rational} \end{matrix}$$


$$\textcircled{2} \quad \left[\frac{r}{s} \right]_q^b = \pm q^{l'} \frac{\overline{\text{hom}}(X, P_2)}{\overline{\text{hom}}(X, P_1)} \quad \begin{matrix} \text{left } q\text{-deformed} \\ \text{rational} \end{matrix}$$

Specialising q_0

Now fix $0 < q < 1$.

Consider the ideal triangle with vertices $0, 1, \infty$.
[corresponds to a piece of stability space]

The $\text{PSL}_2(\mathbb{Z})$ -orbit:




$[q_0 = 1]$

Specialising q

Now fix $0 < q < 1$.

Consider the ideal triangle with vertices $0, 1, \infty$.


The $\text{PSL}_{2,q}(\mathbb{Z})$ -orbit:



[$q = 0.3$]


Specialising \underline{q}

At $q=1$, left & right limits of Farey triangles agree



Specialising g


At $g \neq 1$, the left & right limits of Farey triangles do not agree — we get $[\frac{r}{s}]_g^b$ & $[\frac{r}{s}]_g^\#$!



Specialising g

At $g \neq 1$, the left & right limits of Farey triangles do not agree — we get $[\frac{r}{s}]_g^b$ & $[\frac{r}{s}]_g^{\#}$!

Moreover, the entire semicircle connecting them lies in the limit.



$\overline{\text{Stab}}^q \mathcal{C}$ at a fixed positive q

Thm [B-Becker-Licata]

- ① The union of the closed semicircles $\left[\left[\frac{r}{s} \right]_q^b, \left[\frac{r}{s} \right]_q^\# \right]$ is dense in the boundary of $\overline{\text{Stab}}^q \mathcal{C}$
- ② The remaining points of the boundary are exactly the "q-irrationals".
- ③ The boundary is homeomorphic to S^1 .

Further questions

- Categorical interpretation of q-internals?
- Categorical interpretation of combinatorial properties of left & right q-internals?
- Output from other categories?

Thank you!