

* Last time: Matrix representation of incidence algebra.


Let P be a finite poset.

Choose a labelling (or ordering) of the elements of P

(p_1, p_2, \dots, p_n)

Preferably, choose a topological sorting (matrix looks nicer this way: it is upper-triangular)

Given $f \in \mathcal{A}(P)$, we create a matrix M_f .



** Theorem: Let $f, g \in \mathcal{A}(P)$. Choose an ordering (p_1, \dots, p_n) of P . Let M_f, M_g be the associated matrices of f & g . Then:

(1) The matrix associated to $(f+g)$ is $M_f + M_g$:

$$M_{(f+g)} = M_f + M_g$$


\uparrow addition in $\mathcal{A}(P)$
 \uparrow matrix sum

(2) The matrix associated to $(f * g)$ is $M_f \cdot M_g$:

$$M_{(f * g)} = M_f \cdot M_g$$

\uparrow convolution product
 \uparrow matrix product

** Examples



$f = \zeta$ function

$$g([x, y]) := x + y.$$

$$M_f = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \quad M_g = \begin{bmatrix} 2 & 0 & 4 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}$$

$$(f * g)([x, y]) = \sum_{x \leq z \leq y} f([x, z]) \cdot g([z, y])$$

$$(M_f \cdot M_g)_{(i,j)} = \sum_{1 \leq k \leq n} \underbrace{(M_f)_{(i,k)}}_{f([p_i, p_k]) \text{ or } 0 \text{ if } [p_i, p_k] = \emptyset} \cdot \underbrace{(M_g)_{(k,j)}}_{g([p_k, p_j]) \text{ or } 0 \text{ if } [p_k, p_j] = \emptyset}$$

$$(M_f \cdot M_g)_{(i,j)} = \sum_{\substack{p_i \leq p_k \\ p_k \leq p_j}} \underbrace{f([p_i, p_k])}_x \cdot \underbrace{g([p_k, p_j])}_y$$

Compare w/ previous:


$$(f * g)([x, y]) = \sum_{x \leq z \leq y} f([x, z]) \cdot g([z, y])$$

This comparison proves part (2) of the theorem

$$M_f = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \quad M_g = \begin{bmatrix} 2 & 0 & 4 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}$$

$$(M_f \cdot M_g) = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} & \begin{bmatrix} 2 & 0 & 10 \\ 0 & 4 & 11 \\ 0 & 0 & 6 \end{bmatrix} \end{matrix}$$

it simultaneously gives all values of $(f * g)$ on all intervals.



$$\Rightarrow (f * g)([1, 3]) = 10$$

$$(f * g)([3, 3]) = 6$$

Theorem part (i) is easier:

$$(f + g)([x, y]) = f([x, y]) + g([x, y])$$

$$(M_{f+g})_{(i,j)} = (M_f)_{(i,j)} + (M_g)_{(i,j)}$$

$$= f([P_i, P_j]) + g([P_i, P_j])$$

or zero if $[P_i, P_j] = \emptyset$

$$\Rightarrow M_{f+g} = M_f + M_g$$

** The function μ [Möbius function]

Recall the ζ function in $\mathcal{A}(P)$.

$$(M_\zeta)_{(i,j)} = \begin{cases} 1 & \text{if } [P_i, P_j] \neq \emptyset \\ 0 & \text{otherwise} \end{cases}$$

Suppose we have a topological sorting of P .

(P_1, \dots, P_n)

if $P_i \leq P_j$ then $i \leq j$.

Let $f \in \mathcal{A}(P)$

Suppose that $i > j$. This means that $P_i \not\leq P_j$ [either they are incomparable, or $P_j > P_i$]

In any case, $[P_i, P_j] = \emptyset$.

$$(M_f)_{(i,j)} = 0$$

Given a topological sort, we conclude that all entries below the diagonal in M_f are zero:



Back to ζ : given a topological sort,

$$M_\zeta = \begin{bmatrix} 1 & & & ? \\ 0 & 1 & & \\ 0 & 0 & 0 & \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

1s & 0s depending on which intervals are $\neq \emptyset$.

1s on the diagonal

0s below the diagonal.

From this structure of M_ζ , it is easy to tell whether M_ζ is invertible

** Theorem: An upper-triangular matrix is invertible if and only if all of its diagonal entries are non-zero.

\Rightarrow (Looking at M_ζ): M_ζ is invertible.

$\Rightarrow \zeta$ is always invertible for any poset!

(But how to find it? ...)

** Def: The Möbius function $\mu \in \mathcal{A}(P)$ is the inverse of ζ :

$$\mu * \zeta = \delta = \zeta * \mu$$

i.e. $M_\mu \cdot M_\zeta = \underline{\underline{I}} = M_\zeta \cdot M_\mu$

$$\uparrow \begin{bmatrix} 1 & & 0 \\ & \ddots & \\ 0 & & 1 \end{bmatrix}$$

** Theorem: An element $f \in \mathcal{A}(P)$ is invertible if and only if all diagonal entries of M_f are non-zero.

That is, if and only if $f([x, x]) \neq 0$ for every $x \in P$.