

MATH 2301


* Recap: Matrix representation

Let (P, \leq) be a poset. Fix an ordering (P_1, \dots, P_n) of P .

Given $f \in \mathcal{A}(P)$, we have M_f , an $n \times n$ matrix.

** Key points:

$$- (M_f)_{(i,j)} = \begin{cases} f([P_i, P_j]) & \text{if } [P_i, P_j] \neq \emptyset \\ 0 & \text{otherwise} \end{cases}$$

- If (P_1, \dots, P_n) is a topological sort, then M_f is upper-triangular. 

$$- M_{f+g} = M_f + M_g$$

$$M_{f \times g} = M_f \cdot M_g$$

- $M_{(f^{-1})} = (M_f)^{-1}$ \leftarrow if f is invertible as an element of $\mathcal{A}(P)$.

- Theorem: f is invertible if and only if all diagonal entries of M_f are non-zero.


Equivalently, if and only if $f([x, x]) \neq 0$ for every $x \in P$.

** The Möbius function μ .

Let (P, \leq) be a finite poset.

Def: The Möbius function μ is the inverse of $\zeta \in \mathcal{A}(P)$.

*** Computing μ .



$$\mu * \zeta = \delta.$$


$$(\mu * \zeta)([x, y]) = \delta([x, y]) = \begin{cases} 1 & \text{if } x=y \\ 0 & \text{else.} \end{cases}$$

(1) Case 1: $\mu([x, x]) = ?$ (x some element of P)

$$(\mu * \zeta)([x, x]) = \mu([x, x]) \cdot \underbrace{\zeta([x, x])}_{=1} = \delta([x, x]) = 1$$

↑
from formula

$$\Rightarrow \mu([x, x]) = 1$$



(2) Case 2: $\mu([x, y])$ for $y \neq x$.


E.g. $\mu([a, c])$

$$(\mu * \zeta)([a, c]) = \sum_{a \leq z \leq c} \mu([a, z]) \underbrace{\zeta([z, c])}_{=1} = \delta([a, c]) = 0$$

$$= \mu([a, a]) + \mu([a, c]) = 0 \Rightarrow \mu([a, c]) = -1.$$

Similarly, we have an equation (from $\mu([a, e])$):

$$\underbrace{\mu([a, a])}_{=1} + \underbrace{\mu([a, c])}_{=-1} + \mu([a, e]) = 0 \Rightarrow \mu([a, e]) = 0.$$



$$\mu([a, d])?$$

Equation:

$$\underbrace{\mu([a, a])}_{=1} + \underbrace{\mu([a, b])}_{=-1} + \underbrace{\mu([a, c])}_{=-1} + \mu([a, d]) = 0.$$

$$\Rightarrow \mu([a, d]) = 2 - 1 = 1.$$

In general: suppose $x \neq y$

$$(\mu * \zeta)([x, y]) = \delta([x, y]) = 0.$$


$$\sum_{x \leq z \leq y} \mu([x, z]) \underbrace{\zeta([z, y])}_{=1} = 0.$$

$$\Rightarrow \mu([x, y]) + \sum_{x \leq z < y} \mu([x, z]) = 0.$$

$$\Rightarrow \mu([x, y]) = - \sum_{x \leq z < y} \mu([x, z])$$

recursive formula.

** Example : Divisor poset




Let us compute $\mu([1, -])$

Blue values represent $\mu([1, x])$ at each x .

* There is in fact an intrinsic formula for $\mu([a, b])$ in this case.


** Observe [return to this later]

- If p is a prime, then $\mu([1, p]) = -1$.




- If p, q are different primes, then:

$$\mu([1, pq]) = 1.$$




- If p is a prime, then:

$$\mu([1, p^2]) = 0.$$



** Example : Subset poset

$$S = \{1, 2, 3\}, \quad P(S) = \{A \subseteq S\}$$



Again, let's compute

$$\mu([\emptyset, -])$$

Blue values indicate this

* Again, there is an intrinsic formula for $\mu([A, B])$ in this case.

* Proposition : In a subset poset,

$$(1) \quad \mu([\emptyset, A]) = (-1)^{|A|} \quad \leftarrow \text{number of elements in } A$$

$$(2) \quad \mu([A, B]) = (-1)^{|B \setminus A|} \quad \leftarrow \text{number of elements in the set difference.}$$



** Products of posets

Def: Let (P_1, \leq_1) and (P_2, \leq_2) be posets.

The product poset is the set $P_1 \times P_2$, with the order relation being

$$(a, b) \leq (c, d) \text{ if } a \leq_1 c \text{ and } b \leq_2 d$$

** Examples



$P_1 \times P_2$:

