

MATH 2301

* Nim strategy

** Example

(1, 2, 5, 7)

$\begin{matrix} & 1_2 \\ & 1 0_2 \\ \oplus & 1 0 1_2 \\ \hline & 0 0 1_2 \end{matrix}$
N-position,

according to the theorem.

$$\begin{array}{r} 1_2 \\ 1 0_2 \\ 1 0 1_2 \\ \oplus 1 1 1_2 \\ \hline 0 0 1_2 \neq 0 \end{array}$$

** Theorem

A position in nim is an "N" position iff its nim-sum is non-zero. It is a "P" position iff its nim-sum is zero.

** Proof sketch

We need to show the following:

- 1) Any move from a position with sum = 0 lands us in a position with nonzero sum.
- ✓ 2) From a position with sum $\neq 0$, there is at least one move to a position with sum = 0.

$$\begin{array}{r} 1_2 \\ 1 0_2 \\ 1 0 1_2 \\ \oplus 1 1 1_2 \\ \hline 0 0 1_2 \end{array}$$

Steps to achieve (2):

- 1) Look at first column from the left with an odd number of 1s.
- 2) Choose a pile that has a 1 in that column.

3) Let n be chosen pile, and s be the nim-sum.

4) Take $n \oplus s$, and replace n with $n \oplus s$.

$$\begin{array}{r} 1 0 1_2 \\ \oplus 0 0 1_2 \\ \hline 1 0 0_2 \end{array} \quad \left. \right\} \text{the move } (1, 2, 4, 7) \text{ has nim-sum 0.}$$

More generally: (x_1, \dots, x_k) be a nim config.

$$S = x_1 \oplus \dots \oplus x_k$$

Suppose $S > 0$.

Follow steps ① & ②. Suppose that x_m is a pile size that has a 1 in the leftmost column with an odd number of 1s.

Make the following move: Change x_m to $(x_m \oplus S)$

Why is this a valid move? Is the new nim-sum 0?

$$\begin{array}{r} 1_2 \\ 1 0_2 \\ 1 0 1_2 \\ \oplus 1 1 1_2 \\ \hline 0 0 1_2 \end{array} \quad \text{no} \quad \begin{array}{r} 1_2 \\ 1 0_2 \\ 1 0 0_2 \\ \oplus 1 1 1_2 \\ \hline 0 0 0_2 \end{array}$$

** Prop: The new nim-sum is zero.

Pf: $x_1 \oplus \dots \oplus x_m \oplus \dots \oplus x_k = S$ (old eqn)

$$\begin{aligned} x_1 \oplus \dots \oplus (x_m \oplus S) \oplus \dots \oplus x_k &= ? \quad (\text{new eqn}) \\ \text{commutativity} \quad \text{of } \oplus \quad \swarrow &= S \oplus (x_1 \oplus \dots \oplus x_k) = S \oplus S = 0 \end{aligned}$$

\Rightarrow Prop Under the previous assumptions, $(x_m \oplus s) < x_m$.

Pf:

Note: x_m has a 1 in the first column from the left in which s has a 1.

$$\begin{array}{r} 1_2 \\ 10_2 \\ 101_2 \\ \oplus 111_2 \\ \hline 001_2 \end{array}$$

$x_m \oplus s$ in binary looks like:

$$\begin{array}{r} * * \dots * 1 * * \dots * \\ \oplus 0 0 0 \dots 0 1 * * \dots * \\ \hline * * \dots * 0 * * \dots * \end{array} \quad \left. \begin{array}{l} \rightsquigarrow x_m \} * = \\ \rightsquigarrow s \} \text{unknown.} \end{array} \right.$$

same as something.
 x_m

$$\begin{array}{r} 2 \dots p_1 p \dots 1^0 \\ * * \dots * 1 * * \dots * \rightsquigarrow x_m \text{ contains } 2^p \\ 0 0 0 \dots 0 1 * * \dots * \\ \hline * * \dots * 0 * * \dots * \rightsquigarrow (x_m \oplus s) \text{ does not contain } 2^p, \\ \text{but all higher places equal } x_m. \end{array}$$

This implies that $(x_m \oplus s) < x_m$, because the largest possible number in the second part of $(x_m \oplus s)$ (after the 0) can be $111\dots 1_2 = (1 + 2 + 2^2 + \dots + 2^{p-1})$. This sum equals $(2^p - 1) < 2^p$.

\Rightarrow Changing x_m to $x_m \oplus s$ is a valid nim move.

Part (1) of the proof:

Suppose (x_1, \dots, x_k) is a game position with $x_1 \oplus \dots \oplus x_k = 0$.

Suppose we make a move in x_m , changing it to x'_m .

New nim-sum:

$$x_1 \oplus x_2 \oplus \dots \oplus x'_m \oplus \dots \oplus x_k = s.$$

$$\begin{aligned} \text{Consider } s \oplus 0 &= s = (x_1 \oplus \dots \oplus x'_m \oplus \dots \oplus x_k) \oplus \\ &\quad (x_1 \oplus \dots \oplus x_m \oplus \dots \oplus x_k) \\ s &= x'_m \oplus x_m. \quad (\text{everything else cancels}). \end{aligned}$$

$$\begin{array}{l} x_m = * * \dots * * \\ x'_m = \underbrace{* * \dots * *}_{* * \dots * *} \oplus \\ \qquad\qquad\qquad * * \dots * * = s \end{array} \quad \left. \begin{array}{l} s = 0 \text{ if and only if} \\ x'_m = x_m \text{ binary digit} \\ \text{by digit.} \end{array} \right.$$

So $s = 0 \text{ iff } x'_m = x_m$.


But $x'_m < x_m$ (we made a move)

$$\Rightarrow s \neq 0$$

Grundy labelling

Let G be any impartial combinatorial game.

Eg. : $n = 5$, subtraction game with $S = \{1, 2\}$



Grundy labelling =
more sophisticated
labelling of the game
graph.