

MATH 2301

27/07/2022

* Set theory

* Relations

Def: A relation R on sets S & T is a subset
 $R \subseteq S \times T$

Rule: This is a binary relation (two sets)

In general, you could have multiple sets S_1, S_2, \dots, S_n
& a relation would be a subset $R \subseteq S_1 \times S_2 \times S_3 \times \dots \times S_n$.


Examples

1) ANU MyTimetable enrolment system has lots of relations in the background

such as: $\{ (\text{student}, \text{coursecode}) \mid \text{student is enrolled in the course} \}$

2) $S = \{A, B, C\}$ (names of people)

$T = \{ \text{apple, pear, banana, mango} \}$



Corresponding relation R is:
 $\{ (A, \text{apple}), (A, \text{mango}), (B, \text{pear}), (C, \text{apple}), \dots \}$

Remarks

~~1) Usually, we~~


1) Often in applications, S and T are the same set. In this case, we just say

R is a relation on S .

2) The empty subset $\emptyset \subseteq S \times T$ is a valid relation

3) The full subset $S \times T \subseteq S \times T$ is a valid relation.

* Functions



A function F is a relation on two sets $A \neq B$ such that for every element $a \in A$, there is a unique element $b \in B$ such that $(a, b) \in F$.

We often write this as

$F: A \rightarrow B$, and say $F(a) = b$ if ~~the~~ b is the unique match for $a \in A$.

(3)

Other common properties of relations on a single set S

1) Reflexivity

1) Reflexivity: A relation $R \subseteq S \times S$ is reflexive if $(s, s) \in R$ for every $s \in S$.

2) Symmetry: A relation $R \subseteq S \times S$ is symmetric if whenever $(s, t) \in R$, we also have $(t, s) \in R$.

3) Anti-symmetry: A relation $R \subseteq S \times S$ is anti-symmetric if whenever $(s, t) \in R$ ^{and $s \neq t$,} the pair (t, s) is not in R .

~~[Rmk: this means that pairs such as (s, s) are never in an anti-symmetric relation.]~~ See tomorrow's notes for correction.

4) Transitivity: A relation $R \subseteq S \times S$ is transitive if whenever $(s, t) \in R$ and $(t, u) \in R$, then the pair $(s, u) \in R$.

④

Examples / exercises

$$S = \{a, b, c\}$$

① $R = \{(a, a), (b, b), (c, c)\}$

reflexive ✓
 symmetry ✓
 anti-symmetry ~~✗~~ ✓
 transitivity ✓ (vacuously)

② $R = \emptyset$

reflexivity ✗
 symmetry ✓
 anti-symmetry ✓
 transitivity ✓

} (vacuously)

③ $R = S \times S$

reflexivity ✓
 symmetry ✓
 anti-symmetry ✗
 transitivity ✓

Rmk: We say that an "if x then y " statement is vacuously satisfied if the x part is never satisfied.

Exercise: $S = \mathbb{N} = \{0, 1, 2, 3, \dots\}$

$$R = \{(a, b) \in \mathbb{N} \times \mathbb{N} \mid a \neq 0 \text{ and } b \text{ is divisible by } a\}$$

E.g. $(2, 16) \in R$, but $(3, 19) \notin R$.

Which of the four properties does R have?