

* Midterm Wednesday!

* Extra office hours: 1:30 - 3:00 pm Friday 26th
(Zoom)

2:00 - 3:00 pm Monday 29th
(Hybrid)

Usual on Wed 31st 3:00 - 4:00
(office)

* (P, \leq) a poset

$\mathcal{A}(P) =$ incidence algebra $= \{f: I(P) \rightarrow \mathbb{R}\}$


Convolution product

Inverses w.r.t convolution product:

$f, g \in \mathcal{A}(P)$ are inverses if $f * g = \delta$

Recall: $\zeta \in \mathcal{A}(P)$ is always invertible.

ζ^{-1} is known as $\mu = \mu$ (Möbius function)



* Today: Matrix representations
of $\mathcal{A}(P)$.


Idea: Represent elements of $\mathcal{A}(P)$ as matrices.

Label

Let $f \in \mathcal{A}(P)$

① ~~Make~~ Order the elements of $P = (p_1, \dots, p_n)$
(preferably, in a topological sorting.)

② Make an $n \times n$ matrix [$n = |P|$], rows and
cols indexed by elements of P



$f([p_i, p_j])$ if
 $[p_i, p_j]$ is non-empty.

0 if $[p_i, p_j] = \emptyset$


E.g. (previous example)

$f([x, y]) = \#$ elements in $[x, y]$, if $[x, y] \neq \emptyset$.

matrix:

$$M_f = \begin{matrix} & \begin{matrix} a & b & c & d \end{matrix} \\ \begin{matrix} a \\ b \\ c \\ d \end{matrix} & \begin{bmatrix} 1 & 2 & 2 & 4 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 \end{bmatrix} \end{matrix}$$

Note: this
is upper-triangular



* Proposition: If we choose a topological ordering for the elements of P , then the matrix of any $f \in \mathcal{A}(P)$ will be upper-triangular.

If $f \in \mathcal{A}(P)$, we call the corresponding matrix M_f .

** Addition

$f, g \in \mathcal{A}(P)$

$$(f+g)([x,y]) = f([x,y]) + g([x,y])$$

$$(M_f + M_g)_{(x,y)} = (M_f)_{(x,y)} + (M_g)_{(x,y)}$$

Prop: The matrix of $(f+g)$ is just $M_f + M_g$

$$\text{i.e. } M_{(f+g)} = M_f + M_g$$

** Convolution product


$$(f * g)([x,y]) = \sum_{x \leq z \leq y} f([x,z]) \cdot g([z,y])$$

③

What about

$$(M_f \cdot M_g)_{(x,y)}$$

matrix product.



$$x^{\text{th}} \text{ row of } M_f : ((M_f)_{(x,p_1)}, (M_f)_{(x,p_2)}, \dots, (M_f)_{(x,p_n)})$$

$$y^{\text{th}} \text{ column of } M_g : \begin{pmatrix} (M_g)_{(p_1,y)} \\ (M_g)_{(p_2,y)} \\ \vdots \\ (M_g)_{(p_n,y)} \end{pmatrix}$$

$$(M_f \cdot M_g)_{(x,y)} = \sum_{p_i} \underbrace{(M_f)_{(x,p_i)}}_{=0 \text{ unless } x \leq p_i} \cdot \underbrace{(M_g)_{(p_i,y)}}_{=0 \text{ unless } p_i \leq y}$$

$$(M_f \cdot M_g)_{(x,y)} = \sum_{x \leq z \leq y} (M_f)_{(x,z)} \cdot (M_g)_{(z,y)}$$

④

(5)

** Proposition : The matrix of $(f * g)$ is just

~~of~~ the product $M_f \cdot M_g$, i.e

$$M_{(f * g)} = M_f \cdot M_g$$

\uparrow
convolution
product
 \uparrow
matrix
product

** Proposition : ~~Let~~ Let $f \in A(P)$ be invertible

Then the matrix of f^{-1} is the inverse of
the matrix M_f .

(Move on Monday)