


* Last time: Q: How many integers in $[1, 2023]$ are not divisible by 3, 5, or 7?

We counted:

ints divisible by 3 = $\left\lfloor \frac{2023}{3} \right\rfloor = 674$ (integer part)

ints divisible by 5 = $\left\lfloor \frac{2023}{5} \right\rfloor = 404$

ints divisible by 7 = $\left\lfloor \frac{2023}{7} \right\rfloor = 289$



~~Claim~~
Claim: To find # ints not divisible by any of them, we need to find the # ints divisible by at least one of them & subtract this number from 2023

BUT: If we take $674 + 404 + 289$, that is too many!

Note: The circles overlap, so this has some double-counting!

* Need to add/subtract pairwise, triplewise intersections

ints divisible by 3 and 5: $\left\lfloor \frac{2023}{15} \right\rfloor = 134$

ints divisible by 3 and 7: $\left\lfloor \frac{2023}{21} \right\rfloor = 96$

ints divisible by 5 and 7: $\left\lfloor \frac{2023}{35} \right\rfloor = 57$


ints divisible by 3, 5, and 7: $\left\lfloor \frac{2023}{105} \right\rfloor = 19$
 (triple intersection)

Reminder on principle of inclusion/exclusion:

ints divisible by at least one of 3, 5, 7 is
 $= \sum (\text{single intersections}) - \sum (\text{double intersections}) + \sum (\text{triple intersection})$

$= [674 + 404 + 289] - [134 + 96 + 57] + [19]$

* Consider subset poset of $S = \{3, 5, 7\}$. ③
 $P = \mathcal{P}(S)$



For $A \subseteq S$, let $p(A) = \#$ integers in $[1, 2023]$ that are divisible by everything in A .
 (see orange circled values)

Let $q(A) = \#$ integers in $[1, 2023]$ that are divisible by everything in A and not divisible by anything in $S \setminus A$.

Note: We want $\#$ elts not divisible by anything in S i.e. we want to compute $q(\emptyset)$.

* Note:

$$p(A) = \sum_{A \subseteq B} q(B)$$

E.g. $A = \{3\}$
 $q(A)$ includes things div. by only 3, not 5, not 7
 $q(\{3, 5\}) =$ things div. by only 3 & 5, not 7
 $q(\{3, 7\}) =$ div by 3 & 7, not 5
 $q(\{3, 5, 7\}) =$ div by 3, 5, and 7

Their sum = everything div by 3

(In example):

$$p(\{3\}) = q(\{3\}) + q(\{3, 5\}) + q(\{3, 7\}) + q(\{3, 5, 7\})$$

The previous formula translates to:

$$p = z * q \quad (\text{check using convolution formula})$$

We want the value of q in terms of p :

Use Möbius inversion:

$$q = \mu * p$$

In particular, $q(\emptyset) = (\mu * p)(\emptyset)$.

Formula for μ ?

* Formulas for μ .

① If P is any poset, then:

(a) $\mu([x, x]) = 1$

(b) if $x \not\preceq y$ then

$$\mu([x, y]) = - \sum_{z \in [x, y]} \mu([x, z])$$

$z \in [x, y]$ means $x \preceq z \not\preceq y$.

② If P is a subset poset, then:

(a) $\mu([A, A]) = 1$

(b) If $A \subseteq B$, then

$$\mu([A, B]) = (-1)^{|B \setminus A|}$$

\leftarrow # elts in B that are not in A .

Use this to compute $q(\phi)$ in previous example.

$$q = \mu * p, \quad q(\phi) = \sum_{\phi \subseteq A} \mu([\phi, A]) \cdot p(A)$$

$$q(\phi) = \sum_{\phi \subseteq A} (-1)^{|A|} \cdot p(A)$$

⑤

$$q(\phi) = 2023 - (674 + 404 + 289) + (134 + 96 + 57) - 19$$

$$q(\phi) = 924$$

There are 924 integers in $[1, 2023]$ that are not divisible by 3, 5, or 7.

⑥